



Welfare

of pigs



Victorian Standards and Guidelines

Revision 1

If you would like to receive this information/publication in an accessible format (such as large print or audio) please call the Customer Service Centre on 1300 502 656.

Published by the Department of Primary Industries
Biosecurity Victoria, March 2012.

Published in the Victorian Government Gazette,
8 September 2011

Revision 1 - March 2012

© The State of Victoria 2012.

This publication is copyright. No part may be reproduced by any process except in accordance with the provisions of the *Copyright Act 1968*.

Authorised by the Department of Primary Industries, 1 Spring Street, Melbourne 3000.

ISBN: 978-1-74264-792-0 (print)
ISBN: 978-1-74264-793-7 (online)

Disclaimer

This publication may be of assistance to you but the State of Victoria and its employees do not guarantee that the publication is without flaw of any kind or is wholly appropriate for your particular purposes and therefore disclaims all liability for any error, loss or other consequence which may arise from you relying on any information in this publication.

**For more information about DPI go to
www.dpi.vic.gov.au or phone the Customer
Service Centre on 1300 502 656.**

Victorian Standards and Guidelines for the Welfare of Pigs

Preface

The *Victorian Standards and Guidelines for the Welfare of Pigs* (Pig Welfare Standards) are based on the *Australian Model Code of Practice for the Welfare of Animals – Pigs (3rd edition)* that was prepared by the national Animal Welfare Committee (AWC) and subsequently endorsed by the Primary Industries Ministerial Council on 20 April 2007.

Membership of AWC comprises representatives from each of the State Departments with responsibility for agriculture, CSIRO, Animal Health Australia, and the Australian Government Department of Agriculture, Fisheries and Forestry. Extensive consultation was carried out involving industry, animal welfare organisations and the wider community in the development of the Code of Practice.

Since the development of the *Australian Model Code of Practice for the Welfare of Animals – Pigs*, a new process towards establishing agreed Australian Standards and Guidelines for the Welfare of Animals has commenced. Under the Australian Animal Welfare Strategy, Animal Health Australia has been commissioned to facilitate the development of nationally consistent standards and guidelines, based on the revision of the current Model Codes of Practice for the Welfare of Animals.

The welfare standards and guidelines for livestock aim to streamline livestock welfare legislation in Australia, ensuring that it is both practical for industry and results in improved welfare outcomes. The development of welfare standards and guidelines underpins access to overseas markets and reinforces Australia's international leadership in livestock welfare. Without such change, Australia risks losing consumer confidence and significant national and international markets.

The standards will provide the basis for developing and implementing consistent legislation and enforcement across Australia. The standards are intended for legislation in all jurisdictions.

The process to develop standards both now and in the future involves representatives from livestock industry sectors, animal welfare and research organisations, relevant state and territory government agencies, the Australian Government Department of Agriculture, Fisheries and Forestry and other stakeholders. The process includes a public consultation stage which provides an opportunity for all members of the public to comment on the draft standards and guidelines before they are finalised.

Further information on this process is available at:
www.dpi.vic.gov.au



Contents

Introduction	1
1 Definitions	2
2 Competence of the stock-person	5
3 Food and water	6
3.1 Food	6
3.2 Water	8
4 Accommodation	9
4.1 Accommodation systems	9
4.2 Equipment	14
4.3 Environment	15
4.4 Protection	17
4.5 Waste control	17
4.6 Pigs kept outdoors	18
5 Husbandry	19
5.1 Inspections	19
5.2 Health	20
5.3 Farrowing and weaning	21
5.4 Boar management	22
5.5 Moving pigs	23
5.6 Elective husbandry procedures	24
6 Preparation for transport and slaughter	27
7 Humane destruction (euthanasia)	28
8 Quality assurance systems and record keeping	29

Appendix 1: Condition scoring of pigs	30
Appendix II: Water requirements for pigs	31
Appendix III: Space allowances for pigs	32
Appendix IV: Temperature recommendations	36
Appendix V: Methods for humane destruction (euthanasia) for pigs	37



Introduction

The purpose of the *Victorian Standards and Guidelines for the Welfare of Pigs* (Pig Welfare Standards) is to describe the standards and guidelines that ensure the welfare of pigs in all Australian production systems. The standards and guidelines in this document provide support for assessing and demonstrating high levels of compliance with good farming practices.

The standards (boxed sections) reflect the key management requirements and use the word 'must'. The Standards will be adopted (prescribed) into regulations under the *Livestock Management Act 2010*. The Act and regulations provide for enforcement of those standards.

The standards in the Pig Welfare Standards are based on the knowledge and technology available at the time of publication and will be updated as knowledge and technology evolves. Whilst the document outlines important aspects to be taken into account in ensuring the welfare of pigs, the need for experience and commonsense in the husbandry of animals is also emphasised.

The guidelines are the recommended practices to achieve desirable pig welfare outcomes. Guidelines use the word 'should' and are designed to complement the standards. Non-compliance with one or more guidelines will not constitute an offence under the law.

The Pig Welfare Standards recognise that the basic requirement for the welfare of pigs is a good husbandry system, managed by trained and skilled stock-people.

The specific needs of pigs are:

- readily accessible appropriate and sufficient food and water
- adequate shelter to protect them from climatic extremes
- opportunity to display appropriate patterns of behaviour
- physical handling in a manner which minimises the likelihood of unreasonable or unnecessary pain or distress
- protection from, and/or rapid diagnosis and correct treatment of, injury or disease
- freedom for necessary movement including to stand, stretch, and lie down
- visual and social contact with other pigs.

The Pig Welfare Standards may be used as a reference for inspectors or commercial auditors who are trained and competent to examine and judge the welfare of pigs. Information provided under the headings of 'Guidelines' is advisory only.

1 Definitions

Adult: any pig over the age of nine months.

Ark: a weatherproof moveable structure designed for housing sows and/or piglets in outdoor production systems.

Boar: an uncastrated male pig over nine months of age.

Colostrum: milk secreted by the sow for the first few days after farrowing, characterised by high protein and antibody content.

Condition score: a five-stage scoring system used to classify the condition of pigs, based on the amount of fat and/or muscle covering they have.

Crate: crates are used as independent pieces of equipment and are purpose designed for confining pigs for a number of husbandry functions, including weighing, handling for veterinary interventions, farrowing and assisting with other reproductive processes.

Creep area: a separate area within a farrowing facility in which piglets are protected from crushing or overlying by the sow, and which is usually heated to provide a temperature that is more suitable for maintaining the welfare of piglets, while at the same time maintaining the comfort of the sow.

Deep litter: a type of group housing system in which pigs are kept on a deep layer of bedding material, usually straw or sawdust.

Dry sow: a female pig that has been mated and has not yet farrowed.

Dry sow stall (or gestation sow): an enclosure in which gilts and sows are kept individually. Stalls are normally joined together in rows and may be used for total confinement or allow the pig free choice of access. In addition, the period of confinement may vary from part of the pregnancy to the entire pregnancy.

Elective husbandry procedures: include castration, tail docking, clipping of the needle teeth, nose ringing, identification, back fat measurement, pregnancy diagnosis and tusk trimming.

Farrowing: giving birth to piglets.

Farrowing crate: an enclosure closely related to the sow's body size, in which sows are kept individually during and after farrowing.

Farrowing pen: an enclosure for confining individual sows and their litters during and after farrowing. Such pens contain a creep area and a farrowing crate or other structure for confinement of the sow.

Feeder: equipment from which feed is dispensed.

Feeding stall: an enclosure or stall into which animals enter one at a time to be fed individually.

Finisher: pigs generally above 50 kg live-weight, until they are sold or retained for breeding. The same meaning applies for pigs referred to as 'finishing'. The term 'finisher' usually refers to pigs that are in the final phases of their growth cycle and may include pigs from 50-70 kg.

Foster: a management practice whereby a piglet is moved soon after farrowing, so that it is fed by a sow that is not its mother.

Gestation: the period when a sow is pregnant.

Gestation stall: see dry sow stall.

Gilt: a young female pig, selected for reproductive purposes, before she has been mated.

Grower: pigs generally with live-weights between 20 and 60 kg. The same term can apply for pigs referred to as 'growing' (i.e. throughout the entire growth period cycle from weaning to finishing.)

Growing pigs: weaners, growers and finishers.

Habitually biting: a dog that bites an animal routinely, regularly or constantly.

Health treatment: any medication administered by oral dosing, injection, topical application to the skin or any other means.

Herd health program: a structured and documented management program to identify potential health risks to the pigs and take relevant action to prevent or minimise those risks.

Husbandry: care and management practices in pig keeping.

Hut: see ark.

Lactating sow: a sow that has given birth, and is producing milk to feed her piglets.

Lux: an international measure of light intensity (not to be confused with watts).

Mated gilt: a young female pig that has been mated, but has not had a first litter.

Operator: a person or entity that manages a pig production unit.

Pen: an enclosure for confining pigs in which they can turn around, which may be used for housing pigs in groups, housing boars individually, management purposes such as mating or farrowing, or for confining pigs individually.

Persistent bullying: persistent aggression imposed upon a pig by one or more other pigs.

Piglet: a pig up to the time it is weaned from the sow.

Pork Industry Stockperson Skill Set (PISS): units of competence include:

- care for health and welfare of pigs
- move and handle pigs
- observe enterprise quality assurance procedures, or comply with industry quality assurance requirements
- contribute to OHS processes
- administer medication to animals
- implement animal health control programs
- comply with animal industry requirements.

Reproductive cycle: the period from mating to the following mating, which in the context of this document is defined as 150 days.

Rooting: a behaviour of pigs whereby they use their nose to dig in the ground or in any available material.

Sow: an adult female pig that has had one or more litters. This includes a mated gilt that is confirmed pregnant but yet to give birth.

Special care: includes an individual nutritional regime required for a limited period to restore health and body condition or administration of health treatments to the sow that require her to be confined, such as treatment of injuries and poor body condition resulting from bullying by herd mates.

Stall: an enclosure, closely related to the pig's body size, in which gilts, sows and boars are kept individually. Stalls are normally joined together in rows and may be used for total confinement or allow the pig free choice of access.

Stock-person: a person who undertakes the immediate day-to-day husbandry tasks associated with looking after pigs.

Stock handling: putting into practice the skills, knowledge, experience, attributes and empathy necessary to manage stock.

Stockmanship: the knowledge and skill of caring for pigs.

Suckling piglet: a piglet between birth and weaning, an unweaned pig.

Suitably qualified:

for stock-persons, this means a person who:

- (i) holds a Certificate III in Agriculture (Pig Production) or an equivalent qualification; or

- (ii) has been assessed by a Registered Training Organisation to have successfully completed the units of competence in the Pork Industry Stockpersons Skill Set (PISS); or

- (iii) has worked for a period of at least 12 months, caring for pigs in a commercial pig establishment where the person received on the job training and experience and can demonstrate he or she is competent in the following skill set:

- moving and handling pigs
- inspecting and assessing the health and well-being of pigs
- carrying out vaccinations, health treatments and elective husbandry procedures
- humane destruction of pigs
- maintaining records of inspections and assessments of pigs, or

- (iv) is a veterinary practitioner.

Tethering: a method of restraining pigs whereby a neck or girth collar is attached to a short length of chain, which is in turn fixed to the floor of the front of a pen.

Thoracic sticking: severing the major blood vessels around the heart by inserting a knife into the thoracic cavity in order to drain the blood from an animal.

Under control: in relation to a dog, means that the dog is fully responsive to the commands of the stock-person at all times while mustering livestock.

Weaner: a pig after it has been weaned from the sow up until approximately 30 kg in live-weight.

Weaning: the act of permanently separating piglets from the sow.

2 Competence of the stock-person

Standards

- | | |
|--|--|
| <p>2.1 Persons responsible for the day-to-day needs of pigs must ensure animals under their control are cared for in accordance with the standards in this document.</p> | <p>2.2 Pigs must be cared for by personnel who are skilled in pig husbandry and are suitably qualified to maintain the health and welfare of animals in accordance with the standards in this document, or are under the direct supervision of such personnel.</p> |
|--|--|

Guidelines

- 2.3 Personnel should be appropriately instructed on how their actions may affect a pig's welfare.
- 2.4 Personnel should undergo formal training and/or be trained on-the-job under the supervision of experienced supervisors within the first six months of employment. Thereafter, training should be conducted on a regular basis.
- 2.5 The recommended level of skill and competency for personnel in a supervisory role is a Certificate III in Agriculture (Pig Production), or equivalent qualification.



3 Food and water

3.1 Food

Standards

- | | |
|---|--|
| <p>3.1.1 Pigs must be provided with daily access to feed that maintains their health and meets their physiological requirements.</p> <p>3.1.2 A stock-person responsible for pigs must take remedial action if persistent bullying is leading to deprivation from food.</p> <p>3.1.3 Automatic feeders must be checked daily.</p> | <p>3.1.4 Weaners must be provided with access to feed at least twice daily.</p> <p>3.1.5 If body condition score of a pig falls below 2 (on a scale of 1-5, see Appendix I), action must be undertaken to improve body condition. If remedial action fails to recover them to a score above 2 they must be culled.</p> |
|---|--|

Guidelines

- | | |
|--|--|
| <p>3.1.6 Automated feeding systems should allow animals access to feed in a manner that minimises intimidation, bullying and aggression from other pigs.</p> <p>3.1.7 Food provided should be fresh, palatable, and free from known gross contaminants, physical or toxic substances and micro-organisms at levels that are known to cause harm.</p> | <p>3.1.8 There should be contingencies to provide an alternative means of obtaining and delivering feed, in case of supply failure or delays in delivery.</p> <p>3.1.9 Boars and pregnant sows should be given some bulky or high fibre feed to satisfy appetite. The feed provided for dry sows needs to satisfy appetite without causing the sow to become over-fat.</p> |
|--|--|

3.1.10 Condition scoring can only be used as a guide to assessing the adequacy of nutrition, healthy and productivity for animal welfare. A guide to condition scoring of pigs is given in Appendix I:

- condition score of grower and finisher animals and boars should be 3 or above
- condition score of breeding sows at farrowing should be 3-3.5
- condition score of breeding sows at weaning should be 2.5 or more.

3.1.11 To assess nutritional adequacy, weight for age along with an assessment of the general state of health are more reliable indicators than condition score, particularly in growing pigs.



3.2 Water

Standards		
3.2.1	Drinking water or another wholesome liquid must be easily available to pigs at all times other than during handling and management procedures, to meet their physiological water requirements.	3.2.2 Automatic watering systems must be checked daily.
		3.2.3 A stock-person responsible for pigs must take remedial action if persistent bullying is restricting access to water.

Guidelines

3.2.4	Medicated water should only be used on competent professional advice, as there is a risk that overuse or mixing of medications, or the medication itself, may cause toxic injury to the pigs.	3.2.8	Where wholesome liquid products are to be used as both a food and water source (e.g. whey), advice should be obtained from a qualified adviser on the suitability of the product for that purpose.
3.2.5	Water provided should be palatable ¹ , and at a temperature that does not inhibit drinking.	3.2.9	The daily consumption of water by a pig will vary according to environmental temperature, feeding regime, diet ingredients and live-weight. Providing drinking water at a temperature below 20°C can assist in ensuring adequate intake during hot conditions, where this is possible. Pigs will adapt to drinking warm water in hot climates. Table 2 in Appendix II shows the typical daily water requirements for various classes of pig.
3.2.6	Drinker allocation per pen or group, drinker design, position and flow rates should be such that water requirements of different classes of pig can be met.	3.2.10	Disinfection of drinking water can prevent introduction of diseases. This should be undertaken only according to the instructions of a qualified adviser.
3.2.7	When a piggery is first established, or a new water source obtained, the water can be tested for mineral content and microbiological contamination, and advice obtained on its suitability for pigs from a suitably qualified testing laboratory and/or suitably qualified adviser.	3.2.11	Some pigs might constantly bully others away from a drinker point and it is suggested that more drinker spaces be provided where this occurs.

¹ Reference: Australian and New Zealand Guidelines for Fresh and Marine Water Quality (ANZECC & ARMCANZ), paper 4 – Vol 3 October 2000, ISBN 09578245 0 5.

4 Accommodation

Construction or redesign of housing should be based on expert advice. Housing should meet animal welfare, environmental, and operator health and safety requirements. Pig housing and facilities should be cross-checked for compliance with the standards of this document.

4.1 Accommodation systems

Pigs are currently raised under systems falling into three main categories:

- Indoor (including single and group housing on solid or slatted floors)
- Deep litter (groups on deep litter in shelters or sheds)
- Outdoor (free range in paddocks with shelter such as arks or huts).

Standards		(b) in respect of which:
4.1.1	Accommodation for pigs must be designed, constructed, maintained and managed in a manner that minimises the risks to pigs from adverse weather, injuries or predators.	(i) no development application has been made under the <i>Planning and Environment Act 1987</i> ; and
4.1.2 (1)	The minimum space requirements set out in subclause (2) apply to any new housing, or a substantial alteration to existing housing -	(ii) the building work begins after the commencement of these standards.
	(a) in respect of which a development application is made under the <i>Planning and Environment Act 1987</i> after the commencement of these standards; or	(2) The minimum space requirements for pigs kept in individual housing are: (a) in the case of a sow kept in a stall - the floor space of the stall is not less than 0.6 metres wide and not less than 2.2 metres long; and (b) in the case of a sow kept in a farrowing pen - the floor space of the pen is not less than 5.6 m ² ; and

- (c) in the case of a sow kept in a farrowing crate adjacent to a creep area (whether or not within a farrowing pen):
 - (i) the floor space of the farrowing crate and creep area, when aggregated, is not less than 3.2 m²; and
 - (ii) the floor space of the farrowing crate (which may include space occupied by a rear anti-crush rail, appropriately placed) is not less than 0.5 metres wide (when measured at no more than 450 millimetres from the floor of the crate) and not less than 2 metres long; and
 - (d) in the case of a boar kept in a stall - the floor space of the stall is not less than 0.7 metres wide and not less than 2.4 metres long; and
 - (e) in the case of a boar kept in a pen - the floor space of the pen is not less than 6 m².
- (3) The minimum space requirements for pigs kept in group housing are:
- (a) for each sow kept in the housing - the floor space of the housing is not less than 1.4 m²; and
 - (b) for each gilt weighing more than 100 kilograms kept in the housing - the floor space is not less than 1m²; and
 - (c) in any other case - the floor space of the housing is not less than the number of square metres calculated in accordance with the following formula: $0.030 \times (P^{0.67})$ where P is the average weight in kilograms of all pigs kept in the housing, as set out in Table 4 in Appendix III.

4.1.3 (1)	Sows and boars accommodated individually in stalls must be able to stand, get up and lie down without being obstructed by the bars and fittings of the stall, to lie with limbs extended, to stretch and to be able to undertake such movements freely.	4.1.4	Stalls and farrowing crates must be designed, constructed, maintained and operated to minimise aggression between pigs and overlying of piglets by sows.
(2)	If the pig is kept in a stall:	4.1.5	From 20 April 2017 a sow must not be confined in a stall for more than six weeks of any gestation period. An exception is for individual sows that are under veterinary advice or special care by a competent stock-person.
(a)	it must be able to stand without simultaneously touching both sides or ends of the stall; and		
(b)	when lying down in the stall, the pig's snout and hindquarters must not simultaneously be touching the opposite ends of the stall; and	4.1.6	Facilities for a lactating sow must allow her:
(c)	any bars along the top of a stall must not touch the pig's back when standing or when it has its head down while feeding or drinking; and	(a)	to stand and lie down without obstruction by the bars or fittings of the crate; and
(d)	the placement of drinkers and/or feed/ water troughs in the stall must be easily accessible to the pig, but must not prevent their ability to stand, stretch and lie down; and	(b)	to give birth to piglets without obstruction, and minimise losses of piglets from crushing, trapping or injuries; and
		(c)	to lie down and extend her limbs freely and position herself so that both sides of her udder are accessible to her piglets; and
		(d)	to access feed and water without obstruction.
(e)	when lying down, any contact with their neighbours in stalls on either side must not result in injury.	4.1.7 (1)	Sows must not be confined in farrowing crates for more than six weeks in any one reproductive cycle.

(2)	Subclause (1) does not apply in emergency or exceptional circumstances including when a sow is required to foster a second litter after her own piglets are weaned.	4.1.8	Where boars are kept constantly in stalls they must be released for use for mating or exercised at least twice per week.
		4.1.9	Tethers must not be used to restrain pigs.
(3)	When a sow is required to foster a second litter of piglets, the sow:	4.1.10	Boars run in groups must be monitored daily and managed to ensure that subordinate boars are not seriously injured or subjected to persistent aggression by other boars.
	(a) must not be required to foster more than that second litter; and		
	(b) must be provided with additional care as may be necessary to maintain the good health and vitality of the sow.		

Guidelines

4.1.11	Pigs grouped in pens require sufficient space for each to sleep, defecate and access feed and water.	4.1.14	Aggression problems can occur in group housing accommodation. This may be managed by adopting some or all of the following measures under the advice of qualified advisers. Careful monitoring is required to ensure that aggression is not made worse or merely postponed:
4.1.12	In combination stall/group systems for dry sows or gilts, bullying and deprivation of food can be a problem and this should be monitored and managed by the stock-person.		<ul style="list-style-type: none"> – culling of over-aggressive lines of pigs – drafting out of subordinate sows – grading for size evenness – keeping groups stable
4.1.13	Where individual quarters are provided for dry sows and gilts the partitions should prevent injury from aggressive behaviour while still enabling them to see each other. Bars of stall partitions should be vertical rather than horizontal.		

4.1.15	Mixing of pigs, especially adult sows and boars, is a major cause of aggression in group housing. This may be reduced at the time of mixing by:	– providing environmental enrichment (e.g. provision of straw or novel objects to manipulate)		– grouping after dark
		– provision of escape areas for bullied sows e.g. partitions, feeding stalls	4.1.16	– using masking odours on all pigs in the group.
		– provision of more drinking points		Floors should be installed and maintained in a way that minimises slipping and the risk of injury and allows pigs to stand normally.
		– feeding systems that minimise competition between pigs for feed at feeding times e.g. trickle feeding, individual feeding stalls for sows	4.1.17	Solid surfaces to which pigs have access should be made of materials that can be readily cleaned and disinfected.
		– optimising pen size, shape/design and group numbers	4.1.18	In deep litter systems litter should be replaced or refreshed at intervals that ensure good hygiene, provide for comfort needs and avoid compromise to welfare from scalding and lameness.
		– minimising the mixing of unfamiliar pigs		
		– castration or immuno-castration of males	4.1.19	The provision of straw or other suitable materials to permit foraging behaviour and provide physical and thermal comfort when lying down is encouraged, provided that this is compatible with drainage, hygiene requirements and climatic conditions.
		– adoption of alternative accommodation systems e.g. outdoor management systems, family pen systems		
		– providing free access to feed, provided that over-fatness can be controlled, or free access to feed at the time of first mixing.	4.1.20	Boars can be kept in compatible groups to permit regular exercise. They can also be kept with groups of gilts or sows provided that persistent bullying or excessive unwanted mounting behaviours do not occur.

4.2 Equipment

Standards		
4.2.1	Mechanical equipment essential to provide the basic feed, water and environmental needs of pigs must be inspected at least once a day and maintained in good working order.	
4.2.2	Equipment to which pigs have access must be designed and maintained to minimise risk of injury to the pigs.	
4.2.3	A risk management system must be in place in case of breakdown of mechanical equipment or delay in delivery for alternative ways of providing feed and water, and to provide environmental needs.	
4.2.4	Sheds with automatically controlled forced-ventilation environmental systems (for example: shutters or fans controlled by temperature sensors) must have a backup system and equipment that may be relied on to ventilate the housing in the event of a	
		power failure or equipment malfunction, in order to maintain the health and welfare of pigs.
	4.2.5	In pig housing that is naturally ventilated by means of automatic ventilation:
		(a) automatic ventilation equipment must be inspected twice daily; or
		(b) have alternative ventilation equipment installed that will function automatically in the event of a failure or malfunction of the automatic ventilation equipment; or
		(c) an alarm system installed that will provide a warning of a failure or malfunction of the automatic ventilation equipment.
	4.2.6	Electrical installations at mains voltage must be inaccessible to pigs and properly earthed.

Guidelines

- 4.2.7 Staff skilled in the correct operation of systems (including backup systems) that have a high degree of control over the environment, including automated or mechanical feed delivery systems, should be available to operate such systems.

4.3 Environment

Standards			
4.3.1	Stock-persons must use lighting that enables inspection of all pigs.	4.3.3	Action must be taken to detect and cool heat distressed pigs.
4.3.2	Ventilation must prevent accumulation of harmful concentrations of gases.		

Guidelines

4.3.4	Equipment to routinely measure and record maximum and minimum air temperatures at pig level should be available in all sheds.	4.3.7	At temperatures above 38°C stock-persons should inspect lactating and gestating sows regularly for signs of heat stroke and cool any animal that is affected e.g. water application followed by increasing airflow, provision of ice blocks as a lick; and subject to environmental requirements, provision of access to wallows or mist sprays for outdoor pigs.
4.3.5	Suckling piglets that are under three weeks of age and weaners should be provided with bedding, insulation or supplementary heating that protects against cold. Optimum comfort ranges are outlined in Appendix IV.	4.3.8	In indoor systems, abrupt temperature fluctuations of greater than 10°C during the day should initiate monitoring for adverse effects on pigs that may require manipulation of heating or water cooling and air movement, to assist pigs to cope or be at their optimum temperature range.
4.3.6	During very hot weather (35°C or more) adult pigs are very susceptible to heat stress, and steps should be taken to alleviate distress and avoid deaths. Pigs may die if loaded for transportation in very hot conditions.	4.3.9	Natural or artificial light of (at least 20 lux) is suggested to be made available at pig level in all buildings for a minimum of nine hours daily.

4.3.10 In completely enclosed houses, the level of air exchanges should provide fresh air for respiration, remove excess heat and waste gases, and minimise the effects of dust and excess moisture for pig and human health. The following are guidelines for safe levels of common pollutants for pigs:

Pollutant	Maximum recommended level
Ammonia	11 ppm
Carbon dioxide	1500 ppm
Carbon monoxide	30 ppm
Hydrogen sulphide	5 ppm
Inhalable particles	0.23 mg/m ³

- 4.3.11 Operators are encouraged to have systems in place to measure concentrations of ammonia in enclosed houses. Monitoring is to be focused on areas of least ventilation.
- 4.3.12 Efficient ventilation is particularly important when effluent is held in storage under slatted floors.
- 4.3.13 Guidelines on managing temperature requirements for pigs are provided in Appendix IV.

4.4 Protection

Standards		
4.4.1	All buildings must have fire prevention measures in place in accordance with the requirements of the controlling authority.	shelters that are difficult to service with fire equipment and the pigs are in one space, there must be gates to open to allow pigs to escape.
4.4.2	Approved fire-fighting equipment must be available to service all pig accommodation with staff trained in its use. In large	4.4.3 Preventative measures must be implemented for protection of pigs from predators.

Guidelines

4.4.4	Annual inspection of electrical systems by a qualified person should be conducted.	4.4.7 Pig housing should be located and managed to be safe from the effects of fires and floods.
4.4.5	Fire alarms should be situated on all housing units and be checked regularly for function.	4.4.8 Firebreaks should be established around pasture for open-range systems and pig sheds where the risk of fire is high.
4.4.6	When planning new buildings, consideration should be given to the use of construction materials with a high fire resistance, and all electrical and fuel installations should be planned and fitted so as to minimise the risk of fire.	4.4.9 There should be a sufficient number of exits to facilitate rapid evacuation of personnel and orderly evacuation of pigs where this is practicable, safe for personnel and time permits.

4.5 Waste control

Standard	
4.5.1	Faeces and urine must not be permitted to accumulate to the stage where there is no clean area for pigs to lie down.

4.6 Pigs kept outdoors

Standards		
4.6.1	Access to shelters in cold weather and shade in hot weather must be provided to all outdoor pigs.	4.6.3 Pigs must not be raised on land that is contaminated with toxins, chemical residues, toxic plants or disease-causing organisms at levels that are known to cause harm.
4.6.2	Feed and watering points must be provided so that all pigs can gain access and obtain their daily physiological requirements.	

Guidelines

4.6.4	Space allowances for shelters and grazing should be provided in accordance with Appendix III.	4.6.6 In regions with high summer temperatures, risk mitigation measures should be put in place to reduce the risk of heat stress. This may include the provision of shade, ventilation and subject to environmental considerations, wallows and water sprinklers.
4.6.5	Huts for farrowing and rearing should provide protection for pigs from the elements and provide an environment where pigs are able to manage their thermal comfort.	4.6.7 Consideration should be given to methods of reducing the build up or effect of disease pathogens by the use of herd health programs that include vaccination, parasite control and regular pasture rotation and spelling.
		4.6.8 Commercial breeds of pigs developed for intensive systems may not be suitable for all outdoor conditions. Consideration might be made to use breeds that are more suited such as Large Black, Tamworth, Wessex Saddleback and Berkshire breeds or hybrids of such breeds.



5 Husbandry

5.1 Inspections

Standard

- 5.1.1

Pigs must be inspected at least once a day to assess the pig's health and well-being by a person who is:

(a)

a suitably qualified person; or
- (b)

acting under the supervision of a suitably qualified person.

Guidelines

- 5.1.2

Personnel should be provided with adequate time for the inspection of pigs and the checking of equipment.
- 5.1.3

Personnel in charge of pigs should be able to recognise early signs of distress or disease so that prompt action is taken or advice sought.
- 5.1.4

More frequent and thorough inspections should be undertaken when there is an increased risk to welfare, for example during hot weather, outbreaks of disease, where behavioural vices are occurring, when farrowing is expected, or when groups of pigs have been recently mixed.
- 5.1.5

When pigs are housed in large groups where it is difficult to see all individuals, inspection should be undertaken whilst moving among the livestock.

5.2 Health

Standards			
5.2.1	Persons responsible for the care of pigs must be competent to recognise the signs of ill health, including behavioural anomalies, in pigs and must take appropriate action when any such signs are observed in pigs under their care.	5.2.6	Animals incapable of moving must be destroyed on location.
		5.2.7	A herd health program must be in place to manage the risk of disease.
		5.2.8	If the person in charge is not able to identify the causes of ill health and correct them, they must seek advice from those with training and experience in such matters.
5.2.2	Weaning must be managed to minimise any negative impact on the health and welfare of the sow and piglets.		
5.2.3	Dead pigs must be removed as soon as practicable.	5.2.9	Vaccinations and other health treatments must be administered to pigs only by suitably qualified persons or by persons under the supervision of a suitably qualified person.
5.2.4	Sick, weak or injured pigs must be treated and, if necessary, isolated or destroyed.		
5.2.5	Pigs with incurable sickness, injury or painful deformity must be given appropriate veterinary care or be destroyed.		

Guidelines

5.2.10	Records of sick animals, deaths, treatment given and response to treatment should be maintained to assist disease investigations.	5.2.11	Behavioural vices such as persistent ear, flank or tail biting should be investigated with the assistance of a suitably qualified adviser to identify the possible environment, feed, management or health factors causing the problem.
--------	---	--------	---

5.2.12	The recommended methods of humane destruction (euthanasia) are described in Appendix V.	5.2.14	Methods of carcase disposal can include incineration, composting or deep burial, subject to the approval of the relevant authorities.
5.2.13	Signs of ill health in pigs may include separation from other pigs, lethargy, refusal to eat, reduced production or fertility, changes in the consistency of faeces, vomiting, skin discolouration, shivering, sneezing, coughing, panting, lameness, swellings on the body or joints and abnormal behaviour.		

5.3 Farrowing and weaning

Standards

5.3.1	All piglets must be checked within 24 hours of birth to ascertain that: <ul style="list-style-type: none"> (a) they are feeding; and (b) to ensure that the piglets have had the opportunity to receive colostrum, or have been provided with an appropriate substitute. 	5.3.2	If a sow dies prior to weaning or piglets are receiving inadequate nutrition, the piglets must be fostered, weaned, hand reared or euthanased.
-------	--	-------	--

Guidelines

5.3.3	Sows should be placed in farrowing quarters before the litter is due, to allow them to become accustomed to their surroundings.	5.3.4	If weaning of pigs under three weeks of age, management and nutrition needs to be of very high standard to prevent piglet mortality and ill-thrift.
-------	---	-------	---

5.4 Boar management

Guidelines

5.4.1	Aggressive adult boars should be housed individually to prevent injury from fighting or be kept in compatible groups.	5.4.5	Boars that are raised together are less likely to fight one another and may be compatible in pairs or small groups.
5.4.2	The floor of the service area should be well maintained and not slippery.	5.4.6	Housing systems that provide boars with more freedom of movement than conventional stalls are encouraged for use, provided that such systems are consistent with management of boar hygiene, and operator health and safety requirements.
5.4.3	Matings should be conducted under the supervision of a competent stock-person to prevent aggressive behaviours and injury to boars, sows or gilts.		
5.4.4	Boars may need to be kept in stalls or individual pens to prevent persistent aggression to other boars.		


5.5 Moving pigs

Standard

- | | |
|--|--|
| <p>5.5.1 Electric prodders must not be used on pigs, except during loading, transport or unloading, and only when:</p> <ul style="list-style-type: none">(i) individual pigs weigh 60 kg (live-weight) or more; and(ii) there is reasonable risk to the safety of the driver or the pig(s); and | <p>5.5.2 When using dogs to move pigs the dog/s must be:</p> <ul style="list-style-type: none">(i) under control of a person at all times; and(ii) muzzled if they habitually bite. |
| <p>(iii) other reasonable action to cause movement has failed.</p> | |

Guidelines

- 5.5.3 Pigs should be moved quietly, ideally by using a backing board or other non-injurious objects, by skilled stock-persons.
- 5.5.4 Design of pig housing and loading facilities should be based on expert advice, to facilitate ease of pig movement and minimise stress on animals.



5.6 Elective husbandry procedures

Introduction

The procedures described in this section may be carried out where necessary. Alternative options that minimise or alleviate pain from elective husbandry procedures, or the avoidance of their use, should be adopted where possible.

Standards			
5.6.1	Elective husbandry procedures must not be carried out except by a person:	5.6.2	A surgical procedure to render a male pig sterile must not be performed on a male pig over the age of 21 days unless the procedure is performed:
	(a) suitably qualified to undertake them; or		(a) under anaesthesia; and
	(b) under the direct supervision of a person suitably qualified to conduct the procedures;		(b) by a registered veterinary practitioner.
	and in accordance with these standards.		

Guidelines

5.6.3	Strict attention should be paid to:		– maintenance of good hygiene, particularly of hypodermic syringes, scalpels and needles and of the site of injections
	– suitability of the area in which the procedure is to be performed		– provision of after-care for the animals.
	– the catching and restraining facilities	5.6.4	Restraint used on pigs should be the minimum necessary to complete the procedure safely and quickly.
	– minimising the duration and amount of restraint, pain and distress		
	– the appropriate selection and maintenance of instruments		

Castration

- 5.6.5 If surgical castration is considered necessary in order for market and consumer requirements to be met, it should be performed by a trained and competent operator.
- 5.6.6 Surgical castration requires use of a sterile sharp implement such as a knife or surgical scalpel, with the animal adequately restrained. Good post-operative drainage of the surgical wound is essential.
- 5.6.7 It is recommended that piglets be castrated after two days of age, after they have established their suckling order, and before seven days of age. When pigs 8–21 days of age are castrated, appropriate and effective restraint is necessary.

Tail docking

- 5.6.8 Tail docking should be avoided wherever possible.
- 5.6.9 Where tail biting is a problem, all aspects of the environment, feeding and management should be investigated to identify the contributing factors so that remedial action can be taken e.g. environmental enrichment with straw or other materials that can be manipulated.

- 5.6.10 Where tail docking is practised as a preventative measure, it should be carried out before pigs are seven days of age.

Clipping of ‘needle’ teeth

- 5.6.11 Qualified advice should be sought to determine if teeth-clipping is necessary. This procedure should not be routinely required.
- 5.6.12 If aggression between littermates or damage to the sow is a problem, this procedure should be carried out within three days of birth. It should only be done where unacceptable injury is occurring to littermates and the sow’s udder.
- 5.6.13 Only the tips (no more than a quarter) of the teeth should be removed.

Nose ringing

- 5.6.14 Nose ringing should be avoided. However, this procedure may need to be performed as a last resort, to prevent adverse effects to the environment, if pigs are kept on pasture.
- 5.6.15 Nose rings should be placed through the cartilage of the top of the snout or the tissues separating the nostrils.

- 5.6.16 Provision of adequate substrate or pasture for chewing can provide for exploratory or foraging behaviour and deter pigs from rooting up ground excessively.

Identification

- 5.6.17 Where it is necessary to mark pigs for permanent identification, the ear may be tattooed, tagged, notched or punched, or the body may be tattooed or a micro-chip implanted.
- 5.6.18 Ear notching should be avoided where possible. Where ear notching is performed, it should be carried out before the piglets are seven days of age.

Back-fat measurement and pregnancy diagnosis

- 5.6.19 The recommended method for pregnancy testing and back-fat measurement is with the use of ultrasonic or other non-invasive equipment.

Tusk trimming

- 5.6.20 Tusk trimming of boars is necessary where injury to humans or animals is likely to occur.
- 5.6.21 Tusk trimming should be conducted using embryotomy wire.
- 5.6.22 The boar should be appropriately restrained and, if necessary, anaesthetised for restraint. Analgesia is not required as the tusk lacks sensory nerves.
- 5.6.23 Tusks should be severed cleanly above the level of the gums without causing damage to other tissues.

6 Preparation for transport and slaughter

Pigs should be prepared and transported in accordance with the Australian Animal Welfare Standards and Guidelines – Land Transport of Livestock as prescribed in regulations under the *Livestock Management Act 2010* as well as any other relevant State legislation.



7 Humane destruction (euthanasia)

Introduction

Previous sections of this document have drawn attention to those circumstances when, for humane reasons, pigs may need to be humanely destroyed (euthanased), e.g. if suffering injury or disease.

Standards		
7.1	The method of destruction must cause a sudden unconsciousness with death occurring when unconscious.	(2) Subclause (1) does not apply in urgent circumstances where the services of a suitably qualified person are not reasonably available and destruction of the pig is necessary in order to prevent undue suffering of the pig.
7.2	(1) Destruction of pigs must be done by a suitably qualified person or person acting under direct supervision of a suitably qualified person.	

Guidelines

7.3	The animal should be quietly handled beforehand to ensure it is not unnecessarily distressed or alarmed.	7.4	The methods suitable for humane destruction are detailed in Appendix V.
-----	--	-----	---

8 Quality assurance systems and record keeping

Guidelines






- | | | | |
|-----|---|-----|--|
| 8.1 | It is strongly recommended that all pig farms be part of an appropriate industry approved quality assurance program that includes animal welfare. | 8.3 | Management and monitoring of animal welfare can be aided by the implementation of an audited on-farm quality assurance system that includes the standards in this document. |
| 8.2 | The maintenance of good records is an integral part of a quality assurance system and good farm management. Accurate records should be kept on the case history and treatment of any diseased or injured pigs. Accurate identification of animals is essential. | 8.4 | Quality assurance systems provide a record of welfare, health and productivity data as well as documented evidence of critical management procedures, staff training and details of corrective actions for adverse events. |



Appendix 1:

Condition scoring of pigs

Table 1. Guidelines for body condition score of pigs

Numerical Score	Pelvic Bones, Tail Head	Loin	Vertebrae	Ribs	
1	Pelvic bones very prominent. Deep cavity around the tail head.	Loin very narrow. Sharp edges on transverse spinal process. Flank very hollow.	Prominent and sharp throughout the length of the backbone.	Individual ribs very prominent.	
2	Pelvic bones obvious but some slight cover. Cavity around tail head.	Loin narrow. Only very slight cover to edge of transverse spinal process. Flank rather hollow.	Prominent.	Rib cage less apparent. Difficult to see individual ribs.	
3	Pelvic bones covered.	Edge of transverse spinal processes covered and rounded.	Visible over the shoulder. Some cover further back.	Covered but can be felt.	
4	Pelvic bones only felt with firm pressure. No cavity around tail.	Edge of transverse spinal processes felt only with firm pressure.	Felt only with firm pressure.	Rib cage not visible. Very difficult to feel any ribs.	
5	Pelvic bones impossible to feel. Root of tail set deep in surrounding fat.	Impossible to feel bones. Flank full and rounded.	Impossible to feel vertebrae.	Not possible to feel ribs.	

Care should be taken when assessing body fat and back cover as these can be less in pigs that are selected for certain conformation and fat cover in specific locations.

Appendix II:

Water requirements for pigs

For planning purposes the following consumption estimates are provided for normal ambient temperatures (i.e. 10°C–25°C).

Table 2. Average water consumption (litres per day)*

Boar or dry sow	12–15
Sow and litter	25–45
Grower pig:	
25 kg	3–5
45 kg	5–7
65 kg	7–9
90 kg	9–12

Flow rates will vary depending on the number of drinking points and care should be taken to ensure adequate pump capacity and supply to maintain flow.

Table 3. Recommended water flow rates and maximum water pressures

Class	Flow rate (litres/ minute)	Maximum pressure (kPa)
Weaners	0.5	85-105
Growers/ Finishers	1.0	140-175
Dry sow	1.0	No limit*
Lactating sow	2.0	No limit*

* Care should be made not to have excessive pressure as water wastage can occur.

Appendix III:

Space allowances for pigs

Introduction

Adequate space allowances require a consideration of group size, pen size, age, breed, temperature, ventilation, and lighting.

The following tables of standards for space allowances must be provided for pigs housed indoors and are based on current scientific knowledge and good practice. All dimensions and measures refer to the clear space provided for pigs inside of rails or partitions.

Pen fixtures such as feeders and waterers can be included in this space for stalls and crates as long as they do not impede movement or cause injury.

The space allowances standards for pigs housed indoors outlined below are to be considered binding on all facilities from 20 April 2012, except where otherwise specifically stated.

1. Standards for growing pigs (weaners, growers, finishers)

The minimum available floor area for weaners, growers and finishers is calculated as $\text{m}^2 \text{ per pig} = 0.030 \times \text{bodyweight}^{0.67}$ (See footnote reference²). This formula applies to indoor pens of all flooring types. Where there is a range of weights in a group, the minimum is based on the average weight of pigs in the group.

2 Spoolder, HAM.; Edwards, SA., Corning, S. Livestock Production Science 64 (2000) 167 - 173

Table 4. Minimum space requirement (m² per pig) for weaners, growers, and finishers (LW = Liveweight)

LW(kg)	m²	LW(kg)	m²	LW(kg)	m²	LW(kg)	m²
1	0.03	31	0.30	61	0.47	91	0.62
2	0.05	32	0.31	62	0.48	92	0.62
3	0.06	33	0.31	63	0.48	93	0.63
4	0.08	34	0.32	64	0.49	94	0.63
5	0.09	35	0.32	65	0.49	95	0.63
6	0.10	36	0.33	66	0.50	96	0.64
7	0.11	37	0.34	67	0.50	97	0.64
8	0.12	38	0.34	68	0.51	98	0.65
9	0.13	39	0.35	69	0.51	99	0.65
10	0.14	40	0.36	70	0.52	100	0.66
11	0.15	41	0.36	71	0.52	101	0.66
12	0.16	42	0.37	72	0.53	102	0.67
13	0.17	43	0.37	73	0.53	103	0.67
14	0.18	44	0.38	74	0.54	104	0.67
15	0.18	45	0.38	75	0.54	105	0.68
16	0.19	46	0.39	76	0.55	106	0.68
17	0.20	47	0.40	77	0.55	107	0.69
18	0.21	48	0.40	78	0.56	108	0.69
19	0.22	49	0.41	79	0.56	109	0.70
20	0.22	50	0.41	80	0.57	110	0.70
21	0.23	51	0.42	81	0.57	111	0.70
22	0.24	52	0.42	82	0.57	112	0.71
23	0.25	53	0.43	83	0.58	113	0.71
24	0.25	54	0.43	84	0.58	114	0.72
25	0.26	55	0.44	85	0.59	115	0.72
26	0.27	56	0.45	86	0.59	116	0.72
27	0.27	57	0.45	87	0.60	117	0.73
28	0.28	58	0.46	88	0.60	118	0.73
29	0.29	59	0.46	89	0.61	119	0.74
30	0.29	60	0.47	90	0.61	120	0.74

Guidelines

Pigs housed for more than two weeks in deep litter systems should be provided with at least 30 % more space per pig than the standards listed for group housing with other flooring and waste management systems. This is required to assist with litter management.

2. Standards for breeding gilts, sows and boars housed indoors

Table 5. Minimum space requirements

Class	Minimum space allowance per adult
Gilts in group housing (mated or selected for breeding and >100 kg LW)	1 m ²
Sows in group housing	1.4 m ²
Adult pigs in individual stalls	
all new installations	
Sows	0.6 m x 2.2 m
Boars	0.7 m x 2.4 m
all stalls	Must provide the outcome based standards of Section 4.
Boars in individual pens (living space only)	6.0 m ²
Sows in farrowing crates	
New farrowing crate installations:	
Crate dimensions	0.5 m x 2 m
Farrowing crate and creep area	3.2 m ²
	<ul style="list-style-type: none"> • The minimum length must be 2 metres. This is the internal measurement, inclusive of feed and water facilities and a rear anti-crush rail placed where required. • The minimum width of 500 mm is to be taken at not more than 450 mm above the floor level. • Where crates are smaller than this, they must only be used for smaller sows to achieve the standards of Section 4.1.3.
All farrowing crates	Must provide the outcome based standards of Section 4.1.3.
Farrowing pen	5.6 m ² per sow

3. Guidelines for outdoor pigs

3.1 Space allowances for outdoor systems

Table 6. Maximum stocking rate recommendations in paddocks

Dry sows	20–25 sows/ha
Lactating sows with piglets	9–14 sows/ha

Feeding facilities are included in this allowance.

The following information is common practice:

Sows kept in groups 300–400 m² per sow

Sows kept in individual paddocks 400–500 m² per sow

3.2 Space allowances for shelter accommodation

Table 7. Minimum space allowance recommendations for shelter accommodation

Dry sows in groups	1.2–1.5 m ² /sow
Lactating sow with piglets	4–6 m ² /sow
Boars	2 m ² /boar



Appendix IV:

Temperature recommendations

Pigs, except the very young, are able to tolerate a wide range of temperatures without detriment to their welfare, provided abrupt temperature changes do not occur.

Observation of a pig's behaviour by a competent stock-person is the most reliable method to assess thermal comfort.

When pigs are too cold they will huddle and change position to conserve heat. They will increase their feed intake.

When pigs get too hot they will begin to pant in an effort to cool down and reduce their food intake.

There are a number of ways to manage thermal comfort for pigs e.g. cooling from increased air movement, water sprays, insulation, bedding and supplementary heating.

Table 8. Ranges of temperature that provide optimum comfort for different classes of pigs at pig level

Piglets – newborn	27–35°C
Piglets – three weeks of age	24–30°C
Farrowing house	16–22°C
Weaners	20–30°C in first week
Growers	15–30°C
Finishers	15–30°C
Sows and boars	15–30°C

Appendix V:

Methods for humane destruction (euthanasia) for pigs

Introduction

Humane destruction (euthanasia) is defined as causing a sudden unconsciousness with death occurring when unconscious and without distress, pain, fear or anxiety.

Key points to consider about euthanasia include:

- human safety – staff must be trained to avoid possible injury to themselves or others
- pig welfare – the method must minimise pain and distress to the pig and other pigs
- practicality – the method must be affordable, easy to learn and repeatable
- suitability – the method must be suited to the size of the pig
- location – the procedure must be done in a safe, quiet and private location.

The euthanasia process can be divided into three stages. First, the pig is physically restrained in a way that minimises pain and distress. This may include placing the animal, if small, into the container in which it will be killed. Larger animals may be restrained using a rope snare or placed in a race to restrict the animal's movements. It is then killed in a quick and painless way. Finally, the pig is checked to ensure it is dead.

Methods of euthanasia

There are various methods of euthanasia, described below. The advantages and disadvantages of each of these methods are summarised in the table on the following page.

Guidelines

(i) Carbon dioxide

Carbon dioxide (CO₂) can cause rapid onset of anaesthesia with subsequent death due to respiratory arrest if a concentration of over 80% can be maintained.

The main disadvantage is that pigs can become distressed if the gas is not correctly applied. They have transient muscle spasms before death. However, this is a physiological response after the onset of anaesthesia rather than an indication of stress. The spasms are less intense in stress gene negative pigs than stress gene positive pigs.

Carbon dioxide is heavier than air. Therefore, when constructing a container for pig euthanasia, the outlet valve should be located at the top so the container can be completely filled with carbon dioxide while air is allowed to escape. For small pigs a rubbish bin or similar container with the inlet and outlet valves installed in the lid plus a plastic bag liner, or a cut off inner tube can be used. After checking that the pigs are dead, the bag containing them can be removed.

Table 9. Advantages and disadvantages of each euthanasia method

Method	Human safety risk	Pig welfare	Skill required	Cost	Class of pig
Carbon dioxide (CO₂)	Moderate. Use in well-ventilated area.	Good. Can cause aversive reactions. Causes respiratory arrest following anaesthesia.	Moderate.	Moderate. Initial cost of equipment, CO ₂ supply.	Pigs less than 30 kg.
Anaesthetic overdose	Low if assistance for restraint is available.	Good. Causes respiratory and cardiac arrest following anaesthesia.	High. Must be performed by a registered veterinary practitioner.	Moderate. Anaesthetic solution.	All classes of pigs.
Gunshot	Moderate to high. Training and gun licence required. Security of firearm.	Good. Correct aim essential.	Moderate to high.	Moderate, initial cost of firearm and ammunition.	Pigs greater than 15 kg.
Penetrating captive bolt	Moderate to high. Training required. Security of captive bolt.	Good. Correct aim essential.	Moderate.	Moderate. Initial cost of captive bolt.	Pigs greater than 15 kg.
Blunt trauma to head	Low.	Good if performed on small pigs with rapid force strong enough for instant death.	Low. Proper training required.	None.	Pigs less than 15 kg.

(ii) Anaesthetic overdose

Anaesthetic overdose depresses the central nervous system causing deep anaesthesia leading to respiratory and cardiac arrest. Veterinarians must perform this procedure as it requires intravenous or intracardiac administration. The drugs involved can only be used by registered veterinary practitioners.

(iii) Gunshot

The most efficient and common way to humanely destroy pigs is by a close-range gunshot to the brain.

There may be legal restrictions on discharging a firearm in certain areas. Police permission may be necessary.

A small calibre firearm is most suitable, to reduce the risk of projectiles exiting the head. A 0.22 calibre magnum rifle is adequate for most pigs if the shot is correctly positioned. A 0.22 calibre rifle should only be used on young pigs. The range should be less than 5 m and the muzzle must not be placed against the animal's head.

The animal must be still and properly restrained. Shooting at a moving animal is extremely hazardous to the animal and to bystanders. Never fire while the animal is moving its head.

There are two methods: temporal and frontal.

For the temporal method, the pig is shot from the side of the head so that the bullet enters the skull at a point midway between the eye and the base of the ear on the same side of the head (position 'a' in Figure 1). The bullet should be directed horizontally into the skull. This method is preferred for adult pigs due to the heavier bone structure of the front of the skull.

For the frontal method, the firearm should be directed at a point midway across the forehead and (particularly for adult pigs) about 2 cm above the level of the eyes (position 'b' in Figure 1). Aim horizontally into the skull.




Figure 1. Temporal (a) and frontal (b) positions

(iv) Captive-bolt stunners

The captive-bolt stunner is safer than a firearm, since a blank cartridge is used. The muzzle is firmly pressed against the animal's skull before firing. It must, however, be assumed that the animal has only been stunned and a follow-up method of ensuring death, such as bleeding out, is required. This requires the severing of blood vessels to induce effective bleeding. It is often necessary to follow up a neck cut with a thoracic stick to sever the larger blood vessels at their origin near the heart, because the blood loss from the neck cut alone is not sufficient to ensure the animal does not regain consciousness.

A captive bolt stunner should NOT be used for the temporal approach outlined above for fire-arms.

The manufacturer's directions should be followed on the most appropriate blank cartridge to use for the size of the pig and for storage conditions for cartridges. Regular maintenance of the captive-bolt stunner is essential for efficient stunning.

Two types of captive-bolt stunner are available. The concussion stunner has a wide mushroom-shaped head that delivers a knockout blow to the skull. The penetrating stunner has a narrow bolt that is driven a short distance into the brain. The penetrating type of captive-bolt stunner is recommended, as it is more reliable at delivering an effective stun in pigs. The concussion stunner (non-penetrating) is not recommended.

(v) Stunning by blunt trauma to the head

Blunt trauma to the head using a hammer or other suitable solid heavy object may be used to render unconscious small and easily controlled piglets (up to three weeks old provided they are less than 15 kg). The blow should be aimed at the centre of the forehead in the position indicated for frontal shooting in Figure 1. The unconscious piglet should then be immediately bled out to ensure death.

Evidence of instant death

The following signs indicate that a pig is dead:

- the standing animal will collapse
- the tongue will hang out and be straight and limp
- when a captive bolt is used the eyes will be wide open with a blank stare
- the animal will not blink or have an eye reflex in response to touch
- there will be no evidence of rhythmic breathing or heartbeat
- there will be no response to a nose pinch
- there will be no vocalisation.

If any signs of life are still present the same procedure must be repeated or an alternative approach must be used to kill the animal in a rapid and humane manner.

